

Application of novel fungicide compounds to control the finger millet blast caused by *Pyricularia grisea*

T S S K Patro, N Anuradha, Y Sandhya Rani, U Triveni, K B Palanna, I K Das,
D Sabina Mary, B. Praveen Kumar and M. Divya

Agricultural Research Station, Acharya N. G. Ranga Agricultural University,
Vizianagaram -535 001, A.P.

ABSTRACT

The field experiment was conducted at Agricultural Research Station, Vizianagaram against finger millet blast caused by *Pyricularia grisea* using different fungicides. The per cent disease intensity of leaf blast ranged from 2.0 to 4.7%, neck blast ranged from 9.3 to 83.7% and finger blast ranged from 10.7 to 85.7%. Among all the treatments, T6 (propiconazole) was proved to be best with least incidence of leaf blast (2.0%), neck blast (9.3%) and finger blast (10.7%) and also recorded highest grain yield (1543.3 kg/ha) and fodder yield (4133.7 kg/ha). Treatments, Tebuconazole+ Trifloxystrobin, Tricyclazole, Tricyclazole+ Mancozeb, Isoprothiolane, Azoxystrobin + Difenconazole, Carbendazim + Mancozeb and Carbendazim were also found superior over control in controlling finger millet blast.

Keywords: Blast, Finger millet, New fungicides and Propiconazole.

Millets are the most important cereals food grain crops, especially grown in arid and semi arid regions of the Asia and Africa. Finger millet (*Eleusine coracana*) is popularly known as ragi. It is one of the major food crop and feed as fodder for cattle especially in tribal belt of India. It is a good source of carbohydrates and thus supplies high amount of energy. It is also rich in sulphur containing amino acids, proteins and also has low glycemic index with high fibre increased health problems, due to changes in lifestyle, have driven people to rethink their food habits and deliberately shift toward nutritional crops, such as small millets (Anuradha *et al.*, 2022). Hence, it is recommended for diabetic patients as it is very effective in controlling blood glucose levels of diabetics. High calcium, high soluble fibre, low fat, high diastatic power of malted grains renders finger millet unique. Consumption of finger millet prevents cholesterol and constipation. However, it is traditionally grown in marginal soil conditions with low inputs. The major constraint in the millet growing regions is blast (*Pyricularia grisea*). Blast pathogen attacks all aerial parts of finger millet plant causing

leaf, neck and finger blast and disease appears on leaf lamina with typical spindle shaped spots. The blast disease in finger millet often results in more than 50% yield losses (Esele, 2002) and it is as high as about 80-90% in endemic areas (Viswanath, 1997). Ramappa *et al.* (2002) recorded upto 70 % finger blast and 50 % neck blast during *kharif*, 2000 in Mandya and Mysore districts. Blast disease is considered as number one in the form of yield loss in Andhra Pradesh, Madhya Pradesh, Haryana, Mysore and Maharashtra reported that the ultimate loss in yield is due to enhanced spikelet sterility and reduction in grain weight and number. The most efficient, feasible, ecofriendly and cheapest way to control the plant diseases is the host plant resistance. Efforts are being made to develop finger millet resistance lines to understand inheritance of resistance to *Pyricularia grisea*. (Patro *et al.*, 2013; Patro and Madhuri, 2014; Patro *et al.*, 2016; Patro *et al.*, 2018). However, in rice blast disease is managed primarily through host plant resistance. As, the pathogen has the ability to develop new pathogenic races leading to breakdown of resistance within few years (Ahn, 1994), attempts

have been made to manage blast disease in different crops using fungicide chemicals (Varier *et al.*, 1993; Lukose *et al.*, 2007; Narayana Swamy *et al.*, 2009; Netam *et al.*, 2014; Pagani *et al.*, 2014). However in situ incorporation of legume green manure crops increases the nutrient uptake, productivity of maize and reduce disease incidence (Sandhya Rani *et al.*, 2022). Similarly in groundnut crop simultaneous selection for stable disease resistant and high yielding groundnut genotypes were identified (Patro *et al.*, 2022). Although, host plant resistance is the most economical and viable disease management strategy to control finger millet blast, screening of varieties with inbuilt genetic resistance is the best means for management of this disease, as the crop is predominantly grown by resource poor farmers who can hardly afford using chemicals for its control (Das *et al.*, 2021). In the absence of blast-resistant cultivars, the disease can be best managed with fungicides. Hence, the present study was planned to evaluate eight fungicides against finger millet blast under *in vivo* conditions.

MATERIALS AND METHODS

The field experiments were conducted at Agricultural Research Station, Vizianagaram for the management of blast disease in finger millet by fungicides. The experiment was laid out in randomized block design (RBD) with three replications at spacing of 22.5×10 cm with 3×3 m plot size. Standard agronomic practices of NPK-50kg, 40kg, 25kg were followed at the time of crop growth period. A susceptible variety VR 708 was used in this experiment by imposing the following treatments (Table 1). First foliar spray of fungicides was given at the time of flowering followed by second spray at 10-15 days after first spray. Observations were recorded for leaf, neck and finger blast separately. Leaf blast severity was recorded on 0- 5 scale (Mackill and Bonman, 1992). Whereas, neck blast and finger blast incidence was recorded by counting the number of infected panicles and fingers from total population (Mackill and Bonman, 1992). Disease severity scoring for leaf blast was recorded at seedling and booting stage, whereas for neck blast and finger blast at the physiological maturity and at harvest. The grain yield was recorded after harvesting of crop from individual plots.

RESULTS AND DISCUSSION

The data presented in Table 2 revealed that all the treatments significantly reduced the blast disease when compared to control. The per cent disease intensity of leaf blast ranged from 2.0 to 4.7%, neck blast ranged from 9.3 to 83.7% and finger blast ranged from 10.7 to 85.7%. Among all the fungicides, Propiconazole was effective in managing the blast disease with least per cent disease incidence of leaf blast (2.0%), neck blast (9.3%) and finger blast (10.7%) followed by Azoxystrobin + Difenconazole with leaf blast (2.7), neck blast (16.0) and finger blast (15.0%). The maximum percent disease incidence was recorded in control with leaf blast (4.7), neck blast (83.7) and finger blast (85.7). Propiconazole recorded the maximum grain yield (15.4 q/ha) and fodder yield (41.3 q/ha) followed by Azoxystrobin + Difenconazole with grain yield (14.0 q/ha) while the minimum grain yield (5.8 q/ha) and fodder yield (22.0 q/ha) was recorded in control. Raj and Pannu (2017) [17] reported that Tricyclazole followed by Propiconazole were superior in managing rice blast. Fungicides showed effective control against blast disease in rice ecosystem (Prajapati *et al.*, 2004; Dutta *et al.*, 2012; Sood and Kapoor, 1997) [16, 2, 21]. Carbendazim and Tricyclazole showed effective control against pearl millet blast under field conditions (Lukose *et al.*, 2007; Joshi and Gohel, 2015) [6, 5]. However, rice blast pathogen isolates showed differential sensitivity to Tricyclazole and Carbendazim (Yuan and Yang, 2003; Mohammad *et al.*, 2011) [24, 8]. Narayana Swamy *et al.* (2009) [9] and Ganesh Naik *et al.* (2012) [4] reported that Tebuconazole+ Trifloxystrobin have also been reported to be effective against rice blast. Sharma *et al.* (2018) [19] reported that blast disease can be effectively managed with three sprays of Tebuconazole + Trifloxystrobin or Propiconazole in pearl millet. The results of earlier workers are also in line with the results obtained in the present investigations. Hence, Propiconazole @ 1ml/l was effective in managing all the three blasts, *i.e.*, the leaf blast, neck blast and finger blast disease under *in vivo* conditions in finger millet.

LITERATURE CITED

Ahn SW 1994. International collaboration on breeding for resistance to rice blast. In: Zeigler, R.S., Leong, S.A., Teng, P.S. (Eds.), Rice blast Disease. *CAB International*, Wallingford, UK. 137–153.

Table: 2 Management of Finger millet blast with new fungicide molecules

S.No.	Treatments	Percent Disease Incidence (%)			Grain yield (kg/ha)	Fodder yield (kg/ha)
		Leaf blast	Neck blast	Finger blast		
1	Tebuconazole 50% + Trifloxystrobin 25 WG	3.7	26	30	897.8	2554.17
2	Tricyclazole 75% WP	3.3	25.7	24	1079.5	3032.8
3	Tricyclazole 75% WP + Mancozeb 62% WP	4	22.3	19.3	1302.5	3226.1
4	Isoprothiolane 40% EC	4.3	44	39.7	801.1	2261.6
5	Azoxystrobin + Difenconazole	2.7	16	15	1403.7	3315.5
6	Propiconazole 25% EC	2	9.3	10.7	1543.3	4133.7
7	Carbendazim + Mancozeb	3.7	17.3	18.7	1306	3706.2
8	Carbendazim 50% WP	4.3	28.3	29	992.8	2748.3
9	Control	4.7	83.7	85.7	580.5	2203.3
	CD (5%)	0.9	5.5	6.31	194.04	377.9
	CV (%)	14.4	10.5	12.06	10.19	7.23

Anuradha N, Patro T S S K, Ashok Singamsetti, Sandhya Rani Y, Triveni U, Nirmala Kumari, Nagappa Govanakoppa, Lakshmi Pathy T and Vilas A Tonapi 2022. Comparative study of AMMI and BLUP based simultaneous selection for grain yield and stability of finger millet (*Eleusine coracana* (L.) Gaertn.) genotypes. *Frontiers in Plant Science section Plant Breeding*, Doi:10.3389/fpls.2021.786-839.

Das I K, Palanna K B, Patro T S S K, Ganapathy K N, Kannababu N, Sunil Kumar and Tonapi V A A 2021. multilocational evaluation of blast resistance in a diverse panel of finger millet in India. *Crop protection*. 139:105401

Dutta D, Saha S, Prasad Ray D and Bag M K 2012. Effect of different active fungicides molecules on the management of rice blast disease. *Int. J. Agric. Environ. Biotechnol.* 5:247-251.

Esele J P 2002. Diseases of finger millet. A global review. *Sorghum and finger millet diseases* edited by Leslie JF, 19-26.

Ganesh Naik R, Gangadhara Naik B, Basavaraja Naik T and Krishna Naika R 2012. Fungicidal management of leaf blast disease in rice. *GJBB*. 1:18-21.

Joshi H D and Gohel N M 2015. Management of blast [*Pyricularia grisea* (Cooke) Sacc.] disease of pearl millet through fungicides. *Bioscan*. 10:1855-1858.

Lukose C M, Kadvani D L and Dangaria C J 2007. Efficacy of fungicides in controlling blast disease of pearl millet. *Indian Phytopathol.* 60:68-71.

Mackill D J and Bonman J M 1992. Inheritance of blast resistance in near- isogenic lines of rice. *Phytopathol.* 82:746- 749.

Mohammad R G, Jagadeeshwar R, Krishna Rao V and Rahman S J 2011. Development of fungicidal resistance in *Pyricularia grisea* inducing rice blast to carbendazim. *Indian J. Plant Protect.* 39:215-218.

Narayana Swamy H, Sannaulla S and Dinesh Kumar M 2009. Evaluation of new fungicides against rice blast in cauvery delta. *Karnataka J Agric. Sci.* 22:450-451.

Netam R S, Tiwari R K S, Bahadur A N and Shankar D 2014. *In vitro* and *in vivo* efficacy of fungicides against *Pyricularia grisea* causing finger millet blast disease. *Int. J Plant Protect.* 7:137-142.

Pagani A P S, Dianese A C and Café-Filho A C 2014. Management of wheat blast with synthetic fungicides, partial resistance and silicate and phosphite minerals. *Phytoparasitica*. 42:609-617.

- Patro T S S K, Anuradha N, Ashok Singamsetti, Sandhya Rani Y and Triveni U. 2022.** Simultaneous selection for stable disease resistant high yielding groundnut genotypes. *Current Science*.28(2): 85-92
- Patro T S S K, Anuradha N, Madhuri J, Suma Y and Soujanya A 2013.** Identification of resistant sources for blast disease in finger millet (*Eleusine coracana* Gaertn.). Varietal Improvement of Small Millets. National seminar on “Recent Advances of Varietal Improvement in Small Millets”. 5-6.
- Patro T S S K and Madhuri J 2014.** Identification of resistant varieties of finger millet for leaf, neck and finger blast. *International Journal of Food, Agriculture and Veterinary Sciences*. 4(2):7-11.
- Patro T S S K, Neeraja B, Sandhya Rani Y, Jyothsna S, Keerthi S and Bansal A 2016.** Reaction of elite finger millet varieties against blast disease incited by *Magnaporthe grisea* in vivo. *Progressive Research an International Journal* 11. (2) :209-212.
- Patro TSSK, Meena A, Divya M and Anuradha N 2018.** Evaluation of finger millet early and medium duration varieties against major diseases. *International Journal of Chemical Studies*. 6(3):2184-2186.
- Prajapati K S, Patel R C and Pathak A R 2004.** Field evaluation of new fungicides against blast of rice. *Pestic. Res. J.* 16:26-28.
- Raj R and Pannu P P S 2017.** Management of rice blast with different fungicides and potassium silicate under *in vitro* and *in vivo* conditions. *Journal of Plant Pathology*. 99(3):707-712.
- Ramappa H K, Ravishankar C R and Prakash P 2002.** Estimation of yield loss and management of blast disease in finger millet (ragi). *Proc. Asian Cong. Mycol. Pl. Path. University of Mysore*, 195.
- Sandhya Rani Y, Jamuna P, Triveni U, Patro, T S S K and Anuradha N 2022.** Effect of in situ incorporation of legume green manure crops on nutrient bioavailability, productivity and uptake of maize. *Journal of Plant Nutrition*, 45 (7): 1004–1016.
- Sharma R, Gate V L and Madhavan S 2018.** Evaluation of fungicides for the management of pearl millet (*Pennisetum glaucum* (L.)) blast caused by *Magnaporthe grisea*. *Crop Protection*. 112:209-213
- Sood G K and Kapoor A S 1997.** Efficacy of new fungicides in the management of rice blast. *Plant Dis. Res.* 12:140- 142.
- Varier M, Maiti D and Shukla V D 1993.** Efficacy of combination of fungicide formulations on management of rice-blast (*Pyricularia oryzae*) in rainfed upland. *Indian J. Agric. Sci.* 63:386-389.
- Vishwanath S, Mantur S G and Channamma K A L 1997.** Recent approaches in the management of finger millet diseases. In Proc. National seminar on small millets. ICAR and TNAU, Coimbatore, 27-30.
- Yuan J and Yang X H 2008.** Rice blast fungi (*Magnaporthe grisea*) sensitivity to isoprothiolane and tricyclazole in Guizhou. *Guizhou Agric. Sci.* 6:011.