

# **Application of Novel Fungicide Compounds to Control the Finger Millet Blast caused by *Pyricularia grisea***

**T S S K Patro, N Anuradha, Y Sandhya Rani, U Triveni, K B Palanna, I K Das, D Sabina Mary, B Praveen Kumar and M Divya**  
Agricultural Research Station, ANGRAU, Vizianagaram

## **ABSTRACT**

The field experiment was conducted at Agricultural Research Station, Vizianagaram against finger millet blast caused by *Pyricularia grisea* using different fungicides. The per cent disease intensity of leaf blast ranged from 2.0 to 4.7%, neck blast ranged from 9.3 to 83.7% and finger blast ranged from 10.7 to 85.7%. Among all the treatments, T6 (propiconazole) was proved to be best with least incidence of leaf blast (2.0%), neck blast (9.3%) and finger blast (10.7%) and also recorded highest grain yield (1543.3 kg/ha) and fodder yield (4133.7 kg/ha). Treatments, Tebuconazole+ Trifloxystrobin, Tricyclazole, Tricyclazole+ Mancozeb, Isoprothiolane, Azoxystrobin + Difenconazole, Carbendazim + Mancozeb and Carbendazim were also found superior over control in controlling finger millet blast.

**Keywords:** *New fungicides, finger millet, propiconazole and blast*