



Growth and Yield Influenced by INM in *rabi* Castor Grown on *Vertic Ustochrepts* of South Gujarat

H M Patel, T U Patel, H H Patel, P S Patel and D D Patel

Department of Agricultural Chemistry and Soil Science, Navsari Agricultural University, Navsari-396 450

ABSTRACT

A field experiment was conducted on Inceptisols of College Farm, Navsari Agricultural University, Navsari, (Gujarat) during *rabi* 2004-2005 to study the effect of Integrated nutrient management (INM) in *rabi* castor cv. GCH-4. Application of 5 t bio compost ha⁻¹ along with 100 per cent RDF significantly increased all the growth and yield attributes viz. plant height, number of branches plant⁻¹, length of spike, number of capsules spikes⁻¹ and number of spikes plant⁻¹.

Key words : Castor, Fertilizers, Growth, INM, Organic manure, Yield.

Oil seed crops have a unique significance in recent era of energy crisis as they play pivotal role in the agricultural industry and export trade of India. Castor holds first position among the non-edible oil seeds and occupies 38 and 36 per cent production and area, respectively in India. The integrated management of plant nutrient resources through the judicious use of organic, inorganic and bio-fertilizers offer an opportunity to sustain agriculture along with minimizing any soil and environmental damage. On the contrary, it has been clearly evidenced that the balanced and integrated nutrient management improves the soil organic matter content as well as soil quality which is an index of better soil health and sustainability of the system (Nambiar and Ghosh, 1989; Katyal, 1992 and Rajendra Prasad, 2000). However, the quantification of growth and yield attributes as well as yield of *rabi* castor under south Gujarat condition has not been studied so far, hence, the present study was planned to assess the impact of integrated nutrient management on growth and yield of *rabi* castor.

MATERIAL AND METHODS

A field experiment was conducted during the *rabi* season 2004-05 at College Farm, NAU, Navsari (Gujarat-India), situated at 21° 07' N latitude and 73° 40' E longitude, with an elevation of 10 m above mean sea-level to study The integrated nutrient management (INM) on growth and yield of castor (GCH-4) in south Gujarat. The experiment was laid out in factorial randomized block design (FRBD) with 16 treatments composed of 4 levels of fertilizers (F₀: No fertilizer, F₁: 50 % RDF, F₂: 75 % RDF, F₃: 100

% RDF) and 3 source of manures (M₀ – No organic manure, M₁ – Farm Yard Manure @ 5 t ha⁻¹, M₂ – Bio compost @ 5 t ha⁻¹, M₃ – Vermi compost @ 2 t ha⁻¹) and it was replicated thrice. The experimental soil was classified under jalalpur series with family name fine, montmorillonitic, isohyperthermic *Vertic Ustochrepts*. The color of dry soil is dark brown and texture was clay and neutral to slightly alkaline in reaction. The experimental soil was low in available N, medium in available P₂O₅ and high in available K₂O with slight problem of excess soluble salts and compactness due to the puddling for *kharif* rice crop. Statistical analysis was carried out as per the methods described by Panse and Sukhatme (1967).

RESULTS AND DISCUSSION

The mean data of growth and yield attributes as well as yield of castor as influenced by different levels of inorganic fertilizers, source of organic manures and their interactions are presented in Table 1. The plant height (142.12 cm) and number of branches plant⁻¹ (15.06) were significantly higher with the application of 100 per cent RDF over control and 50 per cent RDF. It might be due to the fact that application of higher quantity of NP might have accelerated the synthesis of chlorophyll and amino acids which are associated with major photosynthetic processes of plants. The results of the present investigation are in confirmation with those of earlier workers Thadoda *et al.* (1996), Hans *et al.* (2001) and Sree and Reddy (2003). Significantly smaller plant (13.43 cm) and less number of branches (6.78) were recorded when no

Table 1. Effect of INM on growth parameters of castor crop

Treatments	Plant height (cm)				
	M ₀	M ₁	M ₂	M ₃	Mean
F ₀	78.33	102.40	106.60	125.67	103.25
F ₁	120.40	143.53	123.40	113.13	125.12
F ₂	101.67	126.13	168.47	124.27	130.13
F ₃	123.20	128.67	170.33	146.27	142.12
Mean	105.90	125.18	142.20	127.33	
	S. Em. ±		C.D. at 5 %		
M	5.65		16.31		
F	5.65		16.31		
M x F	11.29		32.65		
C. V. %	15.63				
No. of branches plant⁻¹					
	M ₀	M ₁	M ₂	M ₃	Mean
F ₀	6.07	6.73	7.60	6.73	6.78
F ₁	6.53	6.40	8.33	7.67	7.23
F ₂	7.00	9.53	13.13	12.00	10.41
F ₃	9.26	14.60	15.06	14.80	13.43
Mean	7.21	9.31	11.03	10.30	
	S. Em. ±		C.D. at 5 %		
M	0.43		1.26		
F	0.43		1.26		
M x F	0.87		2.53		
C. V. %	16.03				

fertilizers were applied. The data further revealed that application of bio compost @ 5 t ha⁻¹ (M₂) produced identical effects on plant height (142.20 cm) and number of branches plant⁻¹ (11.03) at harvest of the crop. The pronounced effect from bio compost over vermi compost noted might be due to the higher rate of application, poor response of FYM might be due to the poor macro and micronutrient content. Application of bio compost might have created better soil condition due to the higher rate of application leading to better mobilization of bound nutrients and improvement in soil aggregates resulting in better root proliferation and thereby efficient absorption of water and nutrients. Secondly, the bio compost has a material which was predigested and also because of presence of pressmud which might have kept the pH of the soil low and thereby increased the availability of micronutrients. Significantly lowest plant height (105.90 cm) and number of branches plant⁻¹ (7.21) at harvest were recorded when organic

manures were eliminated from the treatment. The interactions arising due to the sources of organic manure and levels of inorganic fertilizer significantly altered the growth parameters at the harvest of castor crop. The interaction M₂F₃ (100 % RDF + bio compost @ 5 t ha⁻¹) resulted taller plants (170.33) and more number of branches plant⁻¹ (15.06). The shorter plants were produced when inorganic fertilizers were not applied along with any source of organic manures except, M₃F₀ (Table 1). The influence might be because of continuous supply of available nutrients from inorganic and organic sources on decomposition and due to the production of bio-active substances. The organic manures might have activated the certain growth characters due to the production of bio-active substances and higher fixation of N in the soil.

A perusal of the results presented in Table 2 indicated that length of spike (41.40 cm), number of spikes plant⁻¹ (9.77), number of capsules spike⁻¹ (41.69) and shelling percentage (63.4) were

Table 2. Effect of INM on yield attributes of castor crop.

Treatments	Length (cm) of main spike					Number of spikes plant ⁻¹				
	M ₀	M ₁	M ₂	M ₃	Mean	M ₀	M ₁	M ₂	M ₃	Mean
F ₀	27.73	26.87	27.13	25.73	26.87	7.13	6.80	6.27	6.80	6.75
F ₁	28.33	28.40	29.13	29.00	28.72	6.60	7.20	7.47	7.67	7.23
F ₂	31.33	30.60	38.87	38.80	34.90	7.07	8.93	14.13	8.93	9.77
F ₃	32.87	44.27	46.87	41.60	41.40	7.47	11.53	8.33	11.73	9.77
Mean	30.07	32.53	35.50	33.78		7.07	8.62	9.05	8.78	
	S. Em. ±		C.D. at 5 %			S. Em. ±		C.D. at 5 %		
(M)	1.01		2.93			0.52		1.51		
(F)	1.01		2.93			0.52		1.51		
M x F	2.03		5.86			1.04		3.01		
C. V. %	10.65					21.56				
	Number of capsules per spike					Shelling percentage				
	M ₀	M ₁	M ₂	M ₃	Mean	M ₀	M ₁	M ₂	M ₃	Mean
F ₀	22.13	21.27	28.47	25.53	24.35	46.6	57.8	60.4	59.5	56.1
F ₁	23.47	30.60	30.73	35.00	29.95	46.3	59.3	58.9	55.5	55.0
F ₂	28.67	34.33	37.87	40.07	35.23	52.5	58.7	59.2	61.9	58.1
F ₃	28.60	45.03	48.47	44.67	41.69	60.8	63.3	66.8	62.8	63.4
Mean	25.72	32.81	36.38	36.32		51.6	59.8	61.4	59.9	
	S. Em. ±		C.D. at 5 %			S. Em. ±		C.D. at 5 %		
(M)	1.10		3.18			1.7		5.1		
(F)	1.10		3.18			1.7		5.1		
M x F	2.20		6.36			3.5		NS		
C. V. %	11.63					10.60				

significantly higher with the application of 100 per cent RDF through inorganic fertilizers (F₃). This might be due to the increase in growth traits which helped to increase in synthesis of carbohydrates which was utilized for the development of crop. The data further indicated that application of organic manures significantly changed the length of spike (35.50 cm), number of spikes plant⁻¹ (9.05), number of capsules spike⁻¹ (36.38) and shelling percentage (61.4) with the application of 5 t bio compost ha⁻¹ (M₂). The interaction between organic manures and inorganic fertilizers (M x F) significantly changed the yield attributes of the crop (Table 2). Significantly higher length (46.87 cm) of spike, number of capsules plant⁻¹ (48.47) and shelling percentage (66.8) were observed when 100 per cent RDF along with 5 t bio compost/ha (M₂F₃) were applied. While higher number of spikes plant⁻¹ (14.13) was noted due to application of 75 per cent RDF along with 5 t bio

compost ha⁻¹ (M₂F₂). The scrutiny of results showed the supremacy of organic manures over its no application (M₀). The different organic manures failed to differentiate themselves as far as the number of spikes plant⁻¹ was concerned.

Castor seed and stalk yield was significantly affected due to the different levels of inorganic fertilizers (Table 3). Application of 100 per cent RDF through inorganic fertilizers (F₃) significantly produced 34 per and 50 per cent higher seeds and stalk yield of castor, respectively over control treatment (F₀). The 50 per cent RDF through inorganic fertilizers failed to increase the seed yield in the experiment might be due to the lack of translocation of these nutrients from vegetative parts to yield contributors or might be due to the short supply of nutrients. Similar results were also reported by Sree and Reddy (2003). Application of bio compost @ 5 t ha⁻¹ produced more seed (1691.2 kg

Table 3. Effect of INM on seed and stalk yield (kg ha⁻¹) of castor.

Treatments	Seed				
	M ₀	M ₁	M ₂	M ₃	Mean
F ₀	1256.7	1241.5	1373.6	1345.3	1304.3
F ₁	1301.1	1339.8	1439.4	1291.6	1343.0
F ₂	1374.0	1584.5	1663.9	1390.7	1503.3
F ₃	1486.6	1731.2	2288.0	1485.3	1747.8
Mean	1354.6	1474.3	1691.2	1378.2	
	S. Em. ±		C.D. at 5 %		
M	52.0		150.3		
F	52.0		150.3		
M x F	104.1		300.6		
C. V. %	12.2				
	Stalk				
	M ₀	M ₁	M ₂	M ₃	Mean
F ₀	1083.8	1095.9	1116.6	1097.6	1098.5
F ₁	1144.2	1294.4	1303.0	1337.5	1269.8
F ₂	1270.2	1427.3	1479.1	1543.7	1430.1
F ₃	1348.1	1651.7	2121.7	1473.7	1648.8
Mean	1211.6	1367.3	1505.1	1363.1	
	S. Em. ±		C.D. at 5 %		
M	43.00		124.1		
F	43.00		124.1		
M x F	85.90		248.2		
C. V. %	10.90				

ha⁻¹) and stalk (1505.1 kg ha⁻¹) yield over control while vermi compost and FYM showed more or less similar yield over control. The results evidenced the supremacy of bio compost (M₂) over vermi compost (M₃), FYM (M₁) and no application (M₀) of organic manures could be because of better nutrient supply through out the cropping period and also might be due to the better aggregate formation in soil. Another reason might be increase in yield attributes such as number and length of spikes and number of capsules per spike due to the application of 5 t bio compost ha⁻¹ over no manure. Application of vermi compost and FYM though produced positive effect over control but, failed to increase the seed yield might be due to the different rate of application and quantity of nutrients supplied through them. Significantly 84 and 49 per cent higher castor seed and stalk yields were registered at 100 per cent RDF along with 5 t bio compost/ha (M₂F₃) over the treatment combinations M₁F₀ and M₀F₀, respectively. This was possible due to the improvement in all the growth and yield attributes which might have

generated a cumulative effect on seed yield. Another reason might be due to the additive effect of bio compost which might be possible because of better soil condition, mineralization of essential plant nutrients, addition of micronutrients in the soil, the enhanced activities of microbes and release of growth stimulants (enzymes) due to the increased activities of microorganisms. Similar results were also reported by Arangarasan *et al.* (1999) and Laxminarayana (2004).

LITERATURE CITED

- Arangarasan V, Palaniappan S P and Chelliah S 1999.** Response of castor (*R. communis*) to *Azospirillum* and *phosphobacteria* inoculation. *Journal of Oilseeds Research* 16 (1): 121-122.
- Hans D, Agrawal D K and Sundaramoorthy S 2001.** Appraisal of four castor varieties on the basis of different fertility levels and seed yield. *Current Agriculture* 25 (1-2): 123-125.

- Katyal J C 1992.** Integrated nutrient supply and management potential and possibilities in balanced fertilizer use for increasing food grains production in Northern state (G Dev Ed.). Potash and Phosphorus institute of Canada India Programme, Gurgoan pp: 227–249.
- Laxminarayana K 2004.** Effect of organic and inorganic manures on yield and nutrient uptake of groundnut, *Arachis hypogaea* L. in Ultisols of Mizoram. *Journal of Oil seeds Research* 21 (2): 280-283.
- Nambiar K K M and Ghosh A B 1989.** Highlights of research on a long term fertilizer experiments in India (1971 – 1982). LIFE Research Bult. 1, ICAR, New Delhi.
- Panse V G and Sukhatme P V 1967.** Statistical methods for agricultural workers, ICRA Pub; New Delhi. pp. 152-165.
- Rajendra Prasad 2000.** Nutrient management strategies for the next decade. *Fertilizer News*, 45 (4): 13-18, 21-25 and 27-28.
- Sree S S P and Reddy B B 2003.** Effect of different levels of nitrogen and phosphorus on growth and yield of summer castor after *kharif* rice. *Journal of Oilseeds Research* 20 (1): 93-95.
- Thadoda N M, Sukadia Malkavia D D and Moradia A M 1996.** Response of castor (*Ricinus communis* L.) GCH-4 to planting geometry and N fertilization under rain fed condition. *Gujarat Agricultural University Research Journal* 21 (2): 85-87.

(Received on 13.04.2010 and revised on 05.05.2010)