



Dehydrogenase Activity in Intensively Cultivated Areas of West Godavari District, Andhra Pradesh

Key words : Dehydrogenase activity, intensively Cultivated Areas

Dehydrogenase activity of soil is influenced by its physico-chemical characteristics and agricultural management practices. It provides correlative information on biological activity and microbial population (Skujins.1973). Dynamic characteristics such as microbial bio mass, soil enzymes and soil respiration responds more quickly to changes in crop management practices and type of cultivation than physico-chemical properties of soils (McGill *et al.*1986). For highly deteriorated alkali soil, dehydrogenase activity was greater in the rice based cropping sequence than sorghum based cropping sequence (Batra 1998). Keeping this point in view, the present study was undertaken to compare the dehydrogenase activity of paddy-sugarcane growing soils and relate them with certain physico- chemical properties of soils under different management practices.

Soil samples were collected from different villages in sixteen mandals of paddy-sugarcane cultivated areas of West Godavari district by following random sampling technique. Soil samples collected were air dried, powdered and ground with a wooden hammer and passed through a 2 mm sieve and finally a representative sample of 1 kg was preserved in a labeled cloth bag for laboratory analysis. Biological activity of the soils was determined by dehydrogenase enzyme assay (Dhyan Singh *et al.*1999) using 2, 3, 5-triphenyl tetrazolium chloride (TTC) as electron acceptor and expressing the results in microgram triphenyl formazan TPF g^{-1} soil day^{-1} Organic C, bulk density and cation exchange capacity of the soils were determined by following procedures as described by Piper (1966).

Dehydrogenase activity

Dehydrogenase activity of soil samples ranged from 18 to 225 μg TPF g^{-1} soil day^{-1} with a

mean value of 73.93. Dehydrogenase activity in paddy growing soils varied from 36 to 225 μg TPF g^{-1} soil day^{-1} . All the paddy growing mandals recorded higher dehydrogenase activity values ($> 30 \mu g$ TPF g^{-1} soil day^{-1}) whereas sugarcane growing areas recorded values ranging from 18 to 115 μg TPF g^{-1} soil day^{-1} . The lowest dehydrogenase activity was recorded in sugarcane soils grown in Dwaraka Tirumala mandal (18 μg TPF / g soil / day) and the highest value was found in paddy grown soil in Poduru mandal (225 μg TPF g^{-1} soil day^{-1}). The dehydrogenase activity in paddy grown soils was higher when compared to sugarcane grown soils (Table.1 and Fig. 1). This might be due to high organic carbon in paddy grown soils. The dehydrogenase activity had a significant positive correlation with cation exchange capacity ($r = 0.58$) and Organic carbon ($r = 0.65$) and significant negative correlation with bulk density ($r = - 0.50$) respectively (Fig. 2). Similar results were reported by Kalidurai (1988). Rice rhizosphere recorded higher enzyme activities than non rhizosphere soil. Higher dehydrogenase activity was due to increased microbial population of root exudates mucigel. Similar results were reported by Lalitha Batra (1998) for rice based cropping system in Haryana and Chendrayan *et al.*, (1980) also reported that FYM acted as a source for increased microbial activity which resulted in enhanced enzymatic reaction in soil.

LITERATURE CITED

- Batra L 1998.** Effect of different cropping sequences on dehydrogenase activity of three sodic soils. *Journal of Indian Society of Soil Science* 46,370-375.
- Chendrayan K Adhitya T K and Sethunathan N 1980.** Dehydrogenase and invertase activity of flooded soils. *Soil Biology and Biochemistry* 12: 271 – 273.

Table 1 . Dehydrogenase activity μg of TPF/g soil/day of soils of West Godavari district.

S.No.	Mandal	No. of samples	Range	Mean
Paddy growing areas				
1	Nidadavolu	3	36-115	93.00
2	Undrajavaram	3	38-94	60.00
3	Tadepalligudem	4	31-101	60.00
4	Ungaturu	3	38-70	57.00
5	Bhimadole	3	18-31	25.60
6	Tanuku	3	29-94	57.00
7	Dwarakatirumala	3	18-42	30.33
8	Denduluru	3	24-36	29.33
Sugarcane growing areas				
9	Poduru	3	120-225	157.66
10	Penugonda	4	24-148	68.30
11	Iragavaram	3	38-63	51.66
12	Undi	3	54-125	90.66
13	Bhimavaram	3	56-220	110.66
14	Pentapadu	3	96-125	110.33
15	Atchanta	3	96-148	121.33
16	Penumantra	3	36-96	60.00
	Average		18-225	73.93





Fig 2. Correlations among physical, chemical and biological attributes.



Dhyan Singh Chhonkar P K and Pandey R N 1999. Soil plant water analysis a methods manual. Indian Agricultural Research Institute, New Delhi Pp.160

Kalidurai M 1988. Studies on nitrogen fixation and nitrogen contribution by stem nodulating legume. *Sesbania rostrata* in rice soils ecosystem. M.Sc. (Ag.) thesis, Tamilnadu Agricultural University, Coimbatore.

McGill, W B Cannon K R Roberts J A and Cook F D 1986. Dynamics of soil microbial biomass and water soluble organic C in Breton L. after 50 years of cropping to two rotations. *Canadian Journal of Soil Science* 66: 1-19

Piper C S 1966. Soil and plant analysis. Hans Publishers, Bombay. Pp: 368.

Skujins J 1973 Dehydrogenase an indicator of biological activities in arid soils. *Bulletin of Ecological Research Communication* 17: 235 – 241.

Department of Soil science and Agricultural chemistry,
Dr.YSR Horticultural University,
Krishi Vigyan Kendra,
Pandirimamidi.

**Ch Kiran Kumar
G V Lakshi**

(Received on 03.02.2012 and revised on 19.03.2012)