

**Genetic Variability, Heritability and Genetic Advance for Yield and Yield Component Characters among Restorer Lines of Pearl millet
[*Pennisetum glaucum* (L.) R. Br.]**

A Radhika Ramya, M Lal Ahamed, D RatnaBabu, Y Ashoka Rani, V Srinivasa Rao and R K Srivastava

Department of Genetic and Plant Breeding, Agricultural College, Bapatla 522 101, Andhra Pradesh

ABSTRACT

An investigation was carried out to study genetic variability, heritability and genetic advance among 182 restorer (R-)lines of pearl millet for twelve characters *viz.*, days to 50% flowering, plant height, ear length, ear diameter, productive tillers per plant, head yield per plant, grain yield per plant, panicle harvest index, fresh stover yield per plant, dry matter yield per plant, 1000 grain weight and grain harvest index. The results revealed that the characters ear length, ear diameter, productive tillers per plant, head yield per plant, grain yield per plant, fresh stover yield per plant, dry matter yield per plant and 1000 grain weight showed high PCV and GCV. High estimates of heritability along with genetic advance (% mean) were observed for plant height, ear length, ear diameter, productive tillers per plant, and 1000 grain weight indicating that the selection for these traits would be more effective.

Key words: Genetic Advance, Heritability, Pearl millet, Restorer (R-) lines, Variability.